



# HEALTHCARE NUTRITION COUNCIL

Improving outcomes through awareness and action

*Submitted via Email to: regulations.gov*

September 17, 2021

Centers for Medicare & Medicaid Services  
Department of Health and Human Services  
Attention: CMS-1753-P  
P.O. Box 8010  
Baltimore, MD 21244-1850

**Re: Medicare Program: Hospital Outpatient Prospective Payment and Ambulatory Surgical Center Payment Systems and Quality Reporting Programs; Price Transparency of Hospital Standard Charges; Radiation Oncology Model; Request for Information on Rural Emergency Hospitals; CMS-1753-P**

Dear Administrator Brooks-LaSure:

The Healthcare Nutrition Council (HNC) is providing comments on the CY 2022 Hospital Outpatient Prospective Payment and Ambulatory Surgical Center Payment Systems and Quality Reporting Programs. HNC is an association representing manufacturers<sup>1</sup> of enteral nutrition (EN) formulas and oral nutrition supplements (ONS), parenteral nutrition (PN) formulas, supplies, and equipment. Our mission is to improve health by advancing policies that address and raise awareness of nutrition and its impact on patient outcomes and healthcare costs. HNC aims to promote nutritional screening, diagnosis, assessment, appropriate and timely clinical nutrition interventions, as well as patient access to specialized nutrition support products and services throughout the continuum of care.

In line with our mission, we are providing comments on several areas of this year's proposed rule relating to access to nutritional therapies. As detailed further below, HNC:

- Strongly recommends the inclusion of nutrition quality measures in the Hospital Outpatient Quality Reporting (OQR) Program.
- Strongly supports the expansion of disparities reporting to the Ambulatory Surgical Center (ASC) setting and the inclusion of a diagnosis of malnutrition as part of this reporting.

## **Summary of the importance of addressing malnutrition.**

It is widely recognized that nutritional status plays a significant role in health outcomes and healthcare costs. Addressing malnutrition is essential to improving overall healthcare and may ultimately reduce the economic burden incurred when caring for the oldest and sickest

---

<sup>1</sup> HNC members are Abbott Nutrition, Nestle Healthcare Nutrition, and Nutricia North America.



# HEALTHCARE NUTRITION COUNCIL

Improving outcomes through awareness and action

Americans. Disease-related malnutrition can manifest in patients across all spectrums of body mass index, ranging from under to overweight individuals, including those presenting with obesity. Malnutrition often is associated with acute and chronic diseases and injury, such as cancer, stroke, infection, trauma, and surgical procedures. Large-scale studies have shown that as many as half of hospitalized patients and 35% to 85% of older long-term care residents are undernourished.<sup>1,2,3,4</sup>

If unaddressed, malnutrition increases the cost of care and likelihood of poor health outcomes, including increased complications, longer hospitalizations, and more readmissions. For example, malnourished patients are more likely to experience complications, such as pneumonia,<sup>5</sup> pressure ulcers,<sup>6</sup> nosocomial infections,<sup>7</sup> and death.<sup>8</sup> In addition, malnutrition is a risk factor for other severe clinical events, such as falls<sup>9</sup> and worse outcomes after surgery or trauma.<sup>10</sup> Falls are especially a concern among individuals considered frail. Malnutrition also has negative impacts on patients with specific chronic diseases and conditions, such as stroke,<sup>11</sup> heart failure,<sup>12</sup> cancer,<sup>13</sup> and COPD.<sup>14</sup> Malnourished patients, as well as patients at risk for malnutrition, have significantly longer hospitalizations than well-nourished patients and patients not at risk for malnutrition.<sup>15</sup>

Additionally, readmission rates, institutionalization, and ongoing healthcare services increase in patients suffering from malnutrition. In particular, disease-related malnutrition is a common reason for patients to be readmitted to hospitals.<sup>16</sup> A study published in *HCUP Statistical Briefs*, developed by the Agency for Healthcare Research and Quality (AHRQ), in 2016 found that malnutrition in U.S. hospitalized patients is associated with a more than 50 percent higher rate of readmission within 30 days, compared to patient stays not associated with malnutrition.<sup>17</sup> In 2021, a draft comparative effectiveness review on malnutrition in hospitalized adults, prepared for AHRQ by the Evidence-based Practice Center, found an association between malnutrition and prolonged hospital stays as well as increased mortality among malnourished patients.<sup>18</sup> Hospitalized patients at risk of malnutrition are also more likely to be discharged to another facility or require ongoing healthcare services after being discharged from the hospital than patients who are not at risk.<sup>19</sup>

Beyond just the effect on utilization and outcomes, malnutrition has an outsized effect on overall cost of care. Malnutrition costs associated with older adults aged 65 years and older who are the most at risk for malnutrition, and largely depending on Medicare, are estimated at \$51.3 billion annually.<sup>20</sup> However, this figure likely underestimates the total burden of disease-related malnutrition given the diagnosis gap in hospitalized patients.<sup>21</sup> In addition, malnourished patients and patients with nutrition-related or metabolic issues are frequently readmitted to the hospital.<sup>22</sup> Further, the average costs per readmission for patients with malnutrition were found to be 26-34 percent higher (\$16,900 to \$17,900) compared to those without malnutrition (\$13,400).<sup>23</sup> A retrospective health economic study found that providing oral nutritional supplements (ONS) to Medicare patients aged 65+ with any primary diagnosis was associated with a 16% reduction in length of stay and a 15.8% cost savings – an average of \$3,079 -- per episode.<sup>24</sup>

Lastly, despite the impact on overall health and the prevalence of malnutrition among hospitalized patients, a patient's nutritional status is often not evaluated or diagnosed in a timely manner. In a recent study conducted by AHRQ using the Healthcare Cost and Utilization

Project database, only about 7 percent of hospitalized patients are diagnosed with malnutrition. The extremely low number of those diagnosed with malnutrition represents a screening and diagnosis gap that needs to be addressed.

## **I. HNC strongly recommends the inclusion of nutrition quality measures in the Hospital Outpatient Quality Reporting (OQR) Program.**

CMS is seeking input on the potential future adoption of measures that address care quality in the hospital outpatient setting. It is extremely important to identify malnourished patients and provide them with appropriate nutritional management to ensure recovery, particularly during an emergency room visit. The emergency room treats a broad population of patients and is often used as a source of primary care for individuals without insurance. A CDC report found uninsured adults were more likely than adults with private health insurance or a public health plan to visit the emergency room.<sup>25</sup> As such, screening for malnutrition in emergency room patients would have a positive impact on the health outcomes of the uninsured patient population, which is often impacted by health disparities and malnutrition prevalence. HNC strongly recommends the inclusion of nutrition quality measures in the Hospital OQR Program, and the inclusion of a diagnosis of malnutrition as part of this reporting.

Recent evidence shows the enormous impact malnutrition has on health outcomes in patients diagnosed with COVID-19.<sup>26</sup> The COVID-19 pandemic has elevated the need to identify malnourished patients and ensure a diagnosis of malnutrition, and related interventions, are included in the patient's care plan, and travel with the patient throughout their continuum of care.

Additionally, outside of a healthcare setting, the economic and social consequences resulting from the pandemic contribute to the risk of food insecurity and malnourishment in the community.<sup>27</sup> Malnutrition is a risk factor for severe clinical events, such as loss of lean body mass and risk of falls, and possibly worse outcomes after surgery or trauma, since proper nutrition is critical for healing and recovery. Care provided through an emergency room visit should address these socioeconomic consequences by including an assessment of nutritional status and connecting malnourished patients to community services and access to food. At an August 25 Congressional Briefing titled Older Adult Nutrition Security: Next Steps for Congress, (recording available [here](#)), Meredith Ponder Whitmire, JD, Policy Director at Defeat Malnutrition Today, discussed the availability of community-based federal nutrition services for older adults, including the Supplemental Nutrition Assistance Program (SNAP), noting the considerable body of evidence that shows SNAP plays a role in improving food security, economic security, health, and dietary intake throughout the lifespan.

Malnutrition is a critical issue, and malnutrition screening in the outpatient setting, in particular the emergency room department, is necessary to ensure at-risk patients are identified and cared for before experiencing worsening, associated outcomes.

## **II. HNC strongly supports the expansion of disparities reporting to the Ambulatory Surgical Center (ASC) setting and the inclusion of a diagnosis of malnutrition as part of this reporting.**



# HEALTHCARE NUTRITION COUNCIL

Improving outcomes through awareness and action

CMS is seeking guidance on the potential for measuring disparities in care provided in the Ambulatory Surgical Center Quality Reporting (ASCQR) Program. To date, CMS has not expanded disparities reporting to the ASC setting, and the agency is particularly interested in learning about social risk factors which may highlight social based disparities in facilities with relatively few individuals who possess social risk factors.

CMS notes that significant and persistent inequities in health outcomes exist in the United States, and the COVID-19 pandemic has emphasized many longstanding health inequities, with higher rates of COVID-19 infection and mortality among black, Latino, and Indigenous and Native American persons relative to white persons. HNC commends CMS for addressing these important and ongoing issues and seeking input on the ways social risk factors influence quality of care and outcomes for ASC patients.

Nutritional status, and by consequence malnutrition, is often influenced by a variety of social determinants of health (SDoH). According to the World Health Organization (WHO), SDoH are “the conditions in which people are born, grow, work, live, and age, and the wider set of forces and systems shaping the conditions of daily life. These forces and systems include economic policies and systems, development agendas, social norms, social policies and political systems.”<sup>28</sup> In many cases SDoH will have a drastic impact on the availability and quality of foods, how those foods can be prepared and consumed, and what foods will be commonly consumed as staple parts of the diet. As a result, SDoH shape a population’s nutritional status and may result in certain populations, such as the elderly, disabled, and the poorest segments of society, becoming malnourished.

A patient coming to an ASC from the community could be malnourished but may never have been diagnosed with malnutrition. Patients receiving care in an ASC should be screened for malnutrition, and a diagnosis of malnutrition and related interventions should be included in the patient’s care plan and travel with the patient throughout their continuum of care. As noted above, malnutrition is a risk factor for worse outcomes after surgery or trauma, since proper nutrition is critical for healing and recovery. Additionally, HNC recommends that CMS incorporate nutritional screenings into patients’ Welcome to Medicare preventive visits and annual visits, and ensure malnourished patients are connected to community services and access to food.

HNC strongly recommends that CMS expand disparities reporting to the ASC setting, and include a diagnosis of malnutrition as part of this reporting. Including a diagnosis of malnutrition in quality reporting for the ASC setting will ensure appropriate identification and nutritional management of malnourished patients, make reporting of health disparities based on social risk factors and race and ethnicity more comprehensive, and address gaps in health equity.

\*\*\*

Malnutrition continues to be a crucial component in reducing hospital-acquired conditions, lowering healthcare costs and improving the health and well-being of vulnerable Medicare beneficiaries. **HNC urges CMS to prioritize policies and initiatives that identify and treat malnutrition, encourage proper nutrition and the development of cost-effective nutrition therapy products, and that ensures access through adequate coverage and payment policies for nutrition therapy products.** HNC stands ready to work with CMS and all



# HEALTHCARE NUTRITION COUNCIL

Improving outcomes through awareness and action

stakeholders to develop these policies as one means to improve the public health system. If you have any questions or would like additional information, please contact Justine Coffey, Healthcare Nutrition Council, at [jcoffey@healthcarenutrition.org](mailto:jcoffey@healthcarenutrition.org) or 202-207-1109.

Sincerely,

A handwritten signature in black ink that reads "Robert Rankin".

Robert Rankin  
Executive Director

- 
- <sup>1</sup> Robinson MK, Trujillo EB, Mogensen KM, et al: Improving nutritional screening of hospitalized patients: The role of prealbumin. *JPEN J Parenter Enteral Nutr.* 2003 27:389-395.
- <sup>2</sup> Chima CS, Barco K, Dewitt MLA, et al: Relationship of nutritional status to length of stay, hospital costs, discharge status of patients hospitalized in the medicine service. *J Am Diet Assoc* 1997 97:975-978.
- <sup>3</sup> Braunschweig C, Gomez S, Sheean PM: Impact of declines in nutritional status on outcomes in adult patients hospitalized for more than 7 days. *J Am Diet Assoc* 2000 100:1316-1322.
- <sup>4</sup> Crogan NL, Pasvogel A: The influence of protein-calorie malnutrition on quality of life in nursing homes. *J Gerontol A Biol Sci Med Sci* 2003 58A(2):159-164.
- <sup>5</sup> Callahan CM, Wolinsky FD. Hospitalization for pneumonia among older adults. *J Gerontol.* 1996; 51A:M276-M282.
- <sup>6</sup> Mechanick JI. Practical aspects of nutritional support for wound-healing patients. *Am J Surg.* 2004;188:52S-56S.
- <sup>7</sup> Schneider SM, Veyres P, Pivot X, et al. Malnutrition is an independent factor associated with nosocomial infections. *Br J Nutr.* 2004; 92:105-111.
- <sup>8</sup> Correia MI, Waitzberg DL. The impact of malnutrition on morbidity, mortality, length of hospital stay and costs evaluated through a multivariate model analysis. *Clin Nutr.* 2003;22:235-239.
- <sup>9</sup> Meijers JMM, Halfens RJG, Neyens JCL, et al. Predicting falls in elderly receiving home care: the role of malnutrition and impaired mobility. *J Nutr Health Aging;* 2012; 16: 654-658.
- <sup>10</sup> Marik PE and Flemmer M. Immunonutrition in the surgical patient. *Minerva Anestesiologica.* 2012; 78: 336-342.
- <sup>11</sup> Davalos A, Ricart W, Gonzalez-Huix F, et al. Effect of malnutrition after acute stroke on clinical outcome. *Stroke.* 1996;27:1028-1032.
- <sup>12</sup> Zapatero A, Barba R, Gonzalez N, et al. Influence of obesity and malnutrition on acute heart failure. *Rev Esp Cardiol.* 2012; 65(5): 421-426.
- <sup>13</sup> Lis CG, Gupta D, Lammersfeld CA, et al. Role of nutritional status in predicting quality of life outcomes in cancer – a systematic review of the epidemiological literature. *Nutr J.* 2012; 11:27: 2-18.
- <sup>14</sup> A.S.P.E.N. Board of Directors and the Clinical Guidelines Task Force. Guidelines for the use of parenteral and enteral nutrition in adult and pediatric patients. *JPEN J Parenter Enteral Nutr.* 2002;26(1suppl):1SA-138SA.
- <sup>15</sup> Chima CS, Barco K, Dewitt ML, et al. Relationship of nutritional status to length of stay, hospital costs, and discharge status of patients hospitalized in the medicine service. *J Am Diet Assoc.* 1997; 97: 975-978.
- <sup>16</sup> Alvarez-Hernandez J, Planas Vila M, Leon-Sanz M, et al. Prevalence and costs of malnutrition in hospitalized patients; the PREDyCES® Study. *Nutr Hosp.* 2012; 27(4): 1049-1059.
- <sup>17</sup> Fingar K, Weiss A, Barrett M, Elixhauser A, Steiner C, Guenter P, and Hise Brown M. All-Cause Readmissions Following Hospital Stays for Patients with Malnutrition, 2013. *HCUP Statistical Brief #218.* 2018. 1-18.
- <sup>18</sup> Authors to be included in final report. Draft Comparative Effectiveness Review, Malnutrition in Hospitalized Adults, Prepared for the Agency for Healthcare Research and Quality, June 3, 2021.
- <sup>19</sup> Zapatero A, Barba R, Gonzalez N, et al. Influence of obesity and malnutrition on acute heart failure. *Rev Esp Cardiol.* 2012; 65(5): 421-426.
- <sup>20</sup> Snider J, et al: Economic burden of community-based disease-associated malnutrition in the United States. *JPEN J Parenter Enteral Nutr.* 2014;38:55-165.
- <sup>21</sup> Snider JT, Linticum MT, Wu Y, et al. Economic burden of community-based disease-associated malnutrition in the United States. *JPEN J Parenter Enteral Nutr.* 2014; 38 (Suppl 2): 77S-85S.
- <sup>22</sup> Braunschweig C, Gomez S, Sheean PM. Impact of declines in nutritional status on outcomes in adult patients hospitalized for more than 7 days. *J Am Diet Assoc.* 2000;100:1316-1322.
- <sup>22</sup> Fingar K, Weiss A, Barrett M, Elixhauser A, Steiner C, Guenter P, and Hise Brown M. All-Cause Readmissions Following Hospital Stays for Patients with Malnutrition, 2013. *HCUP Statistical Brief #218.* 2018. 1-18.
- <sup>24</sup> Thomas DR, Zdrowski CD, Wilson MM, et al. Malnutrition in subacute care. *Am J Clin Nutr.* 2002;75:308-313.



# HEALTHCARE NUTRITION COUNCIL

Improving outcomes through awareness and action

---

<sup>25</sup> Renee M. Gindi, Ph.D., et al. Emergency Room Use among Adults Aged 18-64: Early Release of Estimates from the National Health Interview Survey, January-June 2011. Released May 2012. Retrieved from: [https://www.cdc.gov/nchs/data/nhis/earlyrelease/emergency\\_room\\_use\\_january-june\\_2011.pdf](https://www.cdc.gov/nchs/data/nhis/earlyrelease/emergency_room_use_january-june_2011.pdf).

<sup>26</sup> Deepa Handu, PhD, RDN, LDN, Lisa Moloney, MS, RDN, Mary Rozga, PhD, RDN, and Feon W. Cheng, PhD, MPH, RDN, CHTS-CP. Malnutrition Care During the COVID-19 Pandemic: Considerations for Registered Dietitian Nutritionists. *J Acad Nutr Diet* 2021; 121(5): 979-987.

<sup>27</sup> Id.

<sup>28</sup> World Health Organization. Social Determinants of Health. 2019. Retrieved from [http://www.who.int/social\\_determinants/en/](http://www.who.int/social_determinants/en/)